

Pregnant women and the developing fetuses' unique vulnerabilities to environmental hazards

Fragile Beginnings

Children face unique vulnerabilities to environmental hazards at every stage of life.

Children today face a new set of challenges that were unimaginable just a generation ago. Across the world, climate change and environmental degradation are threatening child survival, health and well-being. A fetus's developing cells, organs and tissues are uniquely vulnerable to damage from environmental hazards.

The Fragile Beginnings briefing note series provides a scientific stocktake of the growing body of research on the unique vulnerabilities in utero and at birth, during infancy and childhood, and throughout adolescence. This note focuses on the impact of environmental hazards on developing fetuses. The specific effects of environmental hazards on a pregnant woman's own health are critical but beyond the scope of this document.

Exposure to environmental hazards during pregnancy can have lifelong consequences on a child's health

- Every child has the right to a healthy start in life. But environmental hazards can cause harm before a child is born or even conceived.
- Environmental hazards can damage the egg or sperm cells long before conception takes place, and the effects can be passed down from generation to generation.
- The fetal period is critical as it shapes a child's lifelong health and development.
- There are critical windows of vulnerability to environmental hazards, particularly during the first trimester, when women might not be aware of their pregnancies.
- Adverse birth outcomes such as miscarriages, stillbirth, preterm births, birth defects and low birthweight have been linked to exposure to environmental hazards.
- A fetus's developing cells, organs and tissues are uniquely vulnerable to damage from environmental hazards.
- The consequences can last a lifetime, including health conditions like obesity, cognitive impairment, neurological disorders, lung disease and cancer which can arise in infancy into late adulthood.
- While all women are at risk, the heaviest burden falls upon women in low- and middle-income countries, especially those living in fragile and conflict-affected situations, deepening inequities between groups.
- Preventing exposure to environmental hazards can ensure that future generations live to their full potential. It can also prevent economic loss to countries, including to health systems.



Examples of environmental hazards and how they affect pregnancy

Climate-related hazards

Malaria and Zika

Malaria infection rates in the first trimester of pregnancy are especially high. The parasites that cause malaria can cause severe anaemia in the pregnant woman, pregnancy loss and maternal death. Malaria can also cause babies to be born prematurely and have low birthweight. Children born after antenatal exposure to malaria may have impaired cognitive function.

While pregnant women are no more likely to acquire Zika virus than other women, the virus can seriously harm the fetus's developing nervous system.

Zika virus infection during pregnancy can be devastating, causing pregnancy loss or resulting in babies being born prematurely. Babies who survive may suffer from congenital Zika syndrome, which includes malformation of the head and eyes and significant developmental delays.

Even babies who do not have obvious defects at birth can still present with cognitive and language neurodevelopmental delay in childhood.

Cholera

Cholera is a severe diarrhoeal infection caused by ingesting food or water contaminated with the bacterium *Vibrio cholerae* and can cause dehydration. When a woman is pregnant, severe dehydration can cause problems for the pregnant woman and fetus, including compromised blood flow to the placenta and harmful levels of acid in the amniotic fluid, which can lead to pregnancy loss.

Extreme heat

During pregnancy, a pregnant woman's body adapts to many metabolic changes and maintains a steady and appropriate temperature. But in conditions of extreme heat, a pregnant woman's ability to regulate a healthy temperature can be overwhelmed.

Extreme heat can contribute to dehydration, which can reduce levels of amniotic fluid that protect the fetus.

Extreme heat is associated with preterm birth. It has also been linked to low birthweight, congenital anomalies and gestational diabetes.

Food insecurity

A pregnant woman is especially vulnerable to food insecurity because she needs more nutritious food (both in terms of quantity and quality) so her body can both support the growing fetus and continue to function properly. Climate hazards can reduce the amounts of nutritious food produced, the availability and the cost.

Undernutrition during pregnancy is linked to babies being born too early and too small. It is also linked to neural tube defects, including spina bifida, where the spine and spinal cord do not form properly. Children born to mothers who are experiencing undernutrition because of food insecurity have been shown to have impaired cognitive development.



Pollution

Air pollution

A pregnant woman can be exposed to higher levels of air pollution than a non-pregnant woman due to changes that occur in a woman's breathing system during pregnancy.

Chemicals in air pollution can directly affect a pregnant woman's lungs and create inflammation in her body. They can also enter a pregnant woman's circulation and then cross into the placenta.

Air pollution can affect the health of the pregnant woman (e.g., diabetes in pregnancy, high blood pressure), the ability for her to carry to term and the weight of the baby. It can cause pregnancy loss early on (miscarriage) or later in the pregnancy (stillbirth).

If a child is born after being exposed to air pollution in the womb, that child is at risk of developing neurodevelopmental disorders or asthma, the most common chronic childhood disease.



Lead

There is no safe level of lead exposure.

A fetus's developing brain is particularly vulnerable to lead.

Lead can compete for entry into a pregnant woman's body with important nutrients such as calcium and iron, which are needed at higher levels during pregnancy to support the growth and development of the fetus. Lead can also be released into the pregnant woman's bloodstream from her bones, which store lead from past exposures.

Antenatal exposure to lead is linked to reduced cognitive and behavioural development in children.

Pesticides

Exposure to pesticides can be harmful to a pregnancy, even if a woman is exposed before she gets pregnant.

Pesticides can contribute to impairment of DNA, which can harm the developing fetus. Antenatal exposure to pesticides is linked to birth defects and childhood cancers including neuroblastoma and leukaemia, the most common childhood cancer. Antenatal pesticide exposure can also negatively affect the development of a child's brain.



Arsenic

Arsenic is a toxic heavy metal that can contaminate drinking water and foods. Arsenic can accumulate at levels up to three times higher in the placenta than in a pregnant woman's body.

Arsenic exposure during pregnancy can cause pregnancy loss or premature birth. It is also associated with cognitive deficits in childhood.

Tobacco smoke

When a pregnant woman smokes or is around someone who smokes, her fetus can be exposed to over 7,000 toxic chemicals. There is no safe level of tobacco smoke.

Tobacco smoke can alter blood flow to the fetus, disrupting the delivery of oxygen and nutrients. Babies exposed to tobacco smoke may be born prematurely, have birth defects or die from sudden infant death syndrome. The effects of antenatal smoke exposure can also contribute to many chronic diseases in childhood.



Environmental exposures before and during pregnancy shape a child's lifelong health and well-being

Technical brief

Critical windows of vulnerability

During preconception, conception and all the months of pregnancy, there are critical windows of time where certain cells, organs and tissues are most susceptible to environmental hazards.¹ Understanding these windows can shed light on why environmental hazards can devastate a pregnancy or cause effects on health and development decades later. Protecting pregnant women from environmental hazards is an important step to ensure the health of future generations.

Preconception: The first exposure window


The health of both parents matters. The health of a sperm and egg which combine to create an embryo is foundational to the health of a pregnancy, the fetus and the baby after birth.² The weeks just before conception are particularly important.³ But the impact of environmental hazards can be traced back several generations, and can occur even before a future parent is born.

The damage from direct exposure to environmental hazards to the reproductive cells of both men and women can happen decades before they conceive a child.⁴ This is because a person's reproductive cells are already present when they are in the womb. It has been shown, for instance, that even when a woman does not smoke during pregnancy, her child is at increased risk of asthma if her own mother (the grandmother) smoked while pregnant with her.⁵

Damage can occur anytime, including before conception, continuing in utero, and persist through childhood and adolescence.⁶ Egg and sperm cells also develop differently, which can affect when the windows of sensitivity to environmental hazards occur and, ultimately, fertility and health impacts.⁷

Changes to DNA in the reproductive cells can also be passed on to generations not directly exposed.⁸ This means that a great-grandchild who did not have any direct exposure to an environmental hazard can still develop an environmentally related health condition, such as obesity.⁹

Intergenerational health impacts of exposure to environmental hazards





1. Increased food and fluid intake

A pregnant woman needs more food and fluids than a non-pregnant woman. This increases a pregnant woman's exposure to toxicants in contaminated food and drinking water. For instance, arsenic, a common contaminant of drinking water,¹² is linked to miscarriage, preterm birth and stillbirth,¹³ as well as cognitive deficits in childhood.¹⁴

An increased need for fluids during pregnancy also puts a woman at risk of dehydration. This can cause low amniotic fluid levels, which limits fetal movement and creates pressure on the fetus from the umbilical cord.¹⁵ Diseases like cholera, which make people lose fluids, make pregnant women especially susceptible to dehydration. Severe dehydration can compromise blood flow to the placenta and cause harmful levels of acid in the amniotic fluid, leading to fetal death.¹⁶ Dehydration during heat waves may also contribute to growth restriction and low birthweight.¹⁷

Pregnant women have increased caloric and nutrient needs. This means they are more at risk of complications from food insecurity. Food insecurity is associated with a variety of adverse outcomes, including preterm birth, low birthweight and neural tube defects including spina bifida.¹⁸ Food insecurity in pregnant women is also associated with impaired cognitive development in children.¹⁹

Consuming foods contaminated with fumonisins (mycotoxins found on corn and in corn flour) is associated with neural tube defects.²⁰ Climate change will increase the number of tropical regions susceptible to fungal growth, and this will likely lead to increased food contamination with fumonisins.

2. Changes to the respiratory system

Hormonal changes during pregnancy drive key changes in a pregnant woman's respiratory system, which can affect the body's response to environmental hazards such as air pollution.²¹ Pregnant women inhale more air per minute, which means they will breathe in more air pollutants. Air pollutants can have a dual negative effect because they can both harm the fetus directly and act systemically by creating inflammation and reactive oxygen species, which are unstable molecules that can damage DNA.²²

Antenatal air pollution exposure is associated with miscarriage,²³ preterm birth,²⁴ low birthweight²⁵ and stillbirth.²⁶ It is also linked to common chronic childhood conditions, including neurodevelopmental disorders²⁷ and asthma.²⁸ Exposure to wildfire smoke is linked to preterm birth and low birthweight.²⁹

Exposure to second-hand smoke, which can contaminate children's environments when tobacco products are burned or when a smoker exhales smoke, contains over 7,000 chemicals, including approximately 70 that can

A closer look: How environmental hazards can change DNA instructions in the body

DNA is the genetic material that contains the instructions for a person to develop, survive and reproduce. Environmental hazards can change both the structure and function of DNA. Changes to the structure of DNA are called mutations. Changes to the function of DNA, where genes can be turned on or off, are more common and are called epigenetic changes.¹⁰

Mutations and epigenetic changes can occur in all cells in a body, including reproductive cells. If a mutation or epigenetic change caused by an environmental hazard takes place in a reproductive cell, the damage can be passed on to the next generation.

Epigenetic changes can occur at any point in a person's life but there are certain windows of increased vulnerability, such as during sperm and egg development, fertilization and the formation of an embryo.¹¹

Pregnancy: The second exposure window

During pregnancy, a woman's body undergoes many changes to accommodate the metabolic and nutritional needs of the growing fetus and prepare for breastfeeding. Some of these changes can increase the vulnerability of a pregnant woman and fetus to environmental hazards. They can also alter key processes in a pregnant woman's body and change the way her body responds to physical stresses, such as extreme heat, putting both the woman and the fetus at risk of harm.

cause cancer.³⁰ Prenatal tobacco smoke exposure can alter blood flow and metabolism in the fetus and cause accumulation of chemicals both in the mother and the fetus.³¹ Maternal smoking or second-hand smoke is linked to stillbirths, preterm births, birth defects and infant deaths.³² The effects of prenatal smoke exposure continue through childhood and adolescence and include obesity, attention-deficit/hyperactivity disorder, mood disorders and childhood cancers.³³

Increased demand for oxygen

Increased intake of oxygen necessary to meet the demands of the fetus can increase the production of reactive oxygen species, which are unstable molecules that can damage DNA.³⁴ Pregnancy alters the body's ability to detoxify these reactive oxygen species, and environmental factors can increase their production.³⁵ When there is an imbalance between the production and detoxification of these species, oxidative stress in cells occurs.³⁶

Increased oxidative stress in pregnancy is linked to adverse effects such as pre-eclampsia. It also is linked to the risk of developing cancer during childhood, and metabolic syndrome, cardiovascular disease and neurological diseases later in life.³⁷

3. Changes in movement of substances in the body

The movement of substances, including chemicals, in a body can change during pregnancy. For example, a pregnant woman's bones will remodel to release calcium for the fetus, which can release lead into the bloodstream at the same time.

Other changes include the movement of fats in the body. Early in pregnancy, the storage of fat increases in a woman's body. Later in pregnancy, fat is broken down to provide energy for the fetus.³⁸ 'Fat-loving' (i.e., lipophilic) toxicants can be stored in this fat and then released into a woman's bloodstream, which can potentially reach the fetus.³⁹ Lipophilic chemicals include an array of harmful persistent pollutants, such as DDT, dioxins and heavy metals.⁴⁰

4. Changes in metabolism

Other metabolic shifts to support fetal growth and development include changes in the metabolism of glucose and proteins. These changes are hormonally controlled and susceptible to endocrine-disrupting chemicals (EDCs) and other toxicants. When metabolism during pregnancy is negatively affected by EDCs, diseases such as gestational diabetes, hypertensive disorders or restricted fetal growth can occur. Miscarriage can also happen.⁴¹

5. Changes in the maternal heart, blood and circulation

During pregnancy, the heart increases the amount of blood it pumps per minute. The total volume of blood,

including the plasma (i.e., the liquid portion of blood) is also increased.⁴² Increased plasma volume is critical to maintain blood flow to the uterus. Low plasma volume is associated with gestational hypertension and other pregnancy complications.⁴³

Red blood cell production also increases in pregnancy. As iron is a key component of these cells, pregnant women require more iron.⁴⁴ Conditions such as food insecurity, which may reduce availability of iron from dietary sources, can lead to iron deficiency and anaemia.

6. Changes in heat regulation

Changes in heat regulation to cope with the metabolic changes related to pregnancy include increased blood volume, dilation of blood vessels in the skin and increased sweating.⁴⁵ If a pregnant woman is exposed to extreme heat, she may not be able to transfer enough heat to the external environment. If this happens, her health and the fetus's health can be affected, with consequences for newborn and child health if the pregnancy goes to term.⁴⁶ Extreme heat experienced during pregnancy is associated with preterm birth and has been linked to gestational diabetes, low birthweight and congenital anomalies.⁴⁷

7. Changes in immune function

During pregnancy, a woman's immune system will effectively suppress itself to enable the fetus to grow.⁴⁸ This increases a woman's susceptibility to certain infectious diseases and the severity of illness.⁴⁹ It also increases the risk of fetal death.⁵⁰ Research shows that pregnant women have an increased incidence of many types of infections, including some mosquito-borne illnesses that are increasing due to climate change, such as Zika virus infection and malaria.⁵¹

Malaria has significant impacts on the health of pregnant women and fetuses. Infections in the first trimester are especially high. The parasites that cause malaria can infect red blood cells and accumulate in the placenta, contributing to adverse pregnancy outcomes, including severe anaemia and death in the mother, miscarriage, stillbirth, preterm birth and low birthweight. Malaria during pregnancy has also been shown to be associated with impaired cognitive development in children.⁵²

While Zika virus symptoms during pregnancy are usually mild and similar to those seen in non-pregnant people, Zika virus infection during pregnancy can cause miscarriage, stillbirth and preterm birth.⁵³ Zika virus can affect stem cells in the fetal nervous system.⁵⁴ Babies who survive may suffer from congenital Zika syndrome, which includes having a severely small head, brain and eye anomalies, congenital contractures, intrauterine growth restriction, seizures and neurodevelopmental delay.⁵⁵ Babies who are born without observable defects at birth can still present with cognitive and language neurodevelopmental delay in childhood.⁵⁶ Zika virus can also be transmitted sexually.



The placenta's dual role: Protector and pathway for pollutants and infections

The placenta is a remarkable organ. It takes on critical functions during pregnancy, acting like the lungs, gastrointestinal tract, kidneys and endocrine glands of the developing fetus.⁵⁷ A healthy placenta is essential to a healthy pregnancy.

While the main role of the placenta is to supply the fetus with oxygen and nutrients, the placenta also controls the adaptations a woman's body makes during pregnancy and acts like a barrier to minimize fetal exposures to toxicants and infectious agents.⁵⁸ Unfortunately, many harmful substances, such as lead, mercury, pesticides and other chemicals, can penetrate the placenta barrier and cross through it, harming the fetus and causing potentially lifelong effects to a child.⁵⁹ Some chemicals also concentrate in the placenta at higher levels than in the pregnant woman. For instance, arsenic can accumulate at levels up to three times higher in the placenta compared to the pregnant woman.⁶⁰

Infectious agents, such as bacteria, parasites and viruses, can also cross the placenta and cause serious congenital infections, preterm birth and stillbirth.⁶¹ For example, the Zika virus and a group of antenatal infections known by the acronym TORCH (*Toxoplasma gondii*, other, rubella virus, cytomegalovirus, herpes simplex virus) can devastate the health of the pregnancy and fetus.

A closer look: How environmental hazards affect placenta development and function

Researchers have recently begun to explore how environmental hazards might affect the development of a normally functioning placenta. The placenta has a large number of hormone receptors, which makes it vulnerable to chemicals that can disrupt the endocrine system, including heavy metals, some pesticides and per- and polyfluoroalkyl substances (PFAS).⁶² Endocrine disruption in the placenta can cause oxidative stress and altered epigenetic programming, gene expression and placental function.⁶³

Importantly, abnormal placenta development and function is associated with pre-eclampsia, a condition that includes high blood pressure and kidney damage in a pregnant woman. Pre-eclampsia is one of the most severe complications of pregnancy and can lead to maternal morbidity as well preterm birth and perinatal death.⁶⁴ In children who survive, it can also lead to neurodevelopmental delay and cardiovascular and metabolic disease later in life.⁶⁵ Antenatal air pollution exposure has been shown to be associated with pre-eclampsia.⁶⁶

Exposure during the three pregnancy trimesters

The first trimester (weeks 1 to 13)

The first two weeks after fertilization: Even before a missed period would signal that a woman is pregnant, environmental hazards can interfere with implantation of the embryo into the womb and cause miscarriage.⁶⁷ Also during this period, the patterns of chemical tags on DNA, which can alter the function of DNA if affected by toxicants, are established.⁶⁸ By the time the embryo is implanted, the majority of DNA methylation tagging patterns are established.⁶⁹

Weeks 4 to 8: Organs begin to form. By the eighth week of pregnancy, all major external and internal structures are formed, and organ systems have started to develop. During this time, the embryo's tissues and organs are rapidly developing, which makes them especially vulnerable to environmental hazards. As a result, weeks 4 to 8 is the window of time when major birth defects can occur.⁷⁰

From week 9: This starts the fetal period. Until birth, organs continue to develop and mature and remain

vulnerable to environmental hazards. Critical windows of development correspond to the periods of time that cells are dividing and changing into different types and tissues and the times that organs take on their shape.

For brain development, the critical period is from weeks 3 to 16. But brain development can continue to be affected by environmental hazards after this time as the brain will continue to grow and develop.


The second trimester (weeks 14 to 27)

This is a time of rapid growth and development. During the second trimester, lungs are most vulnerable to impairment from air pollution.⁷¹

The third trimester (weeks 28 to 40)

This is when fetal weight increases and organs continue to mature so they can be ready to function at birth.⁷² During this time, environmental exposures can alter the functional maturity of organ systems, including the central nervous system.⁷³

Timeline of health impacts from exposure to environmental hazards during trimesters



Note: An array of pregnancy-related disorders, including those caused by environmental hazards, can lead to the death of the embryo (miscarriage).

All women are at risk of environmental hazards during pregnancy – but some women are more at risk than others

Women in low- and middle-income countries, including fragile and humanitarian contexts

Women living in poverty and in low- and middle-income countries are more likely to suffer ill effects from environmental hazards.⁷⁴ A woman's health at the start of pregnancy and her genetic make-up can also play a role. Those particularly at risk include women who live in regions where unclean fuels are still used to cook, heat and light homes, as this generates high levels of household air pollution.⁷⁵ Others particularly at risk are those in regions where contamination of food crops with fumonisins (a group of mycotoxins) is common. Women working in dangerous jobs and without adequate workplace regulations, such as informal e-waste recycling, artisanal small-scale gold mining and in agricultural roles where pesticides are used, are also at great risk

as they are regularly exposed to harmful chemicals.⁷⁶ Women can also be exposed to hazardous chemicals in female-dominated workplaces such as hairdressers and nail salons, or in daily living from cosmetics and household products (such as cleaning agents).⁷⁷

The increased risk to health for pregnant women in urban slums

Living in an urban slum significantly affects the health of a pregnant woman and her developing fetus. Overcrowded and unsanitary conditions increase the prevalence of infectious diseases. Pregnant women in urban slums often experience inadequate nutrition and lack access to quality health care services. Air and water pollution are common, as slums are often located near industrial sites or waste dumps, exposing pregnant women to harmful chemicals.⁷⁸



How environmental hazards affect fetal, newborn and child health

Health outcome	Causes
<p>Numerous health conditions in a pregnant woman can affect the pregnancy</p>	<p>The health status of a pregnant woman is critical to the health of the developing fetus and pregnancy.</p> <p>High blood pressure, gestational diabetes and obesity can contribute to adverse pregnancy outcomes and can be affected by environmental chemicals. For example, phthalates (chemicals added to plastics and found in many personal care and household products), PFAS, polychlorinated biphenyls (PCBs, a persistent organic pollutant formerly used in the electricity industry) and some flame retardants (such as polybrominated diphenyl ethers or PBDEs) significantly increase the risk of gestational diabetes.⁷⁹ Air pollution is also associated with gestational diabetes and high blood pressure.⁸⁰</p>
<p>Death</p>	<p>Exposure to environmental hazards is associated with an array of pregnancy-related disorders that can lead to miscarriage, stillbirth and sudden infant death syndrome.⁸¹</p>
<p>Preterm birth</p>	<p>Many environmental hazards are linked to preterm birth (i.e., being born before 37 weeks of gestation).⁸²</p> <p>Some examples include air pollution, lead, some pesticides, PFAS and phthalates.⁸³</p> <p>Complications from preterm births are the leading cause of death among children under 5 years of age. Being born premature is also associated with many developmental problems in childhood.⁸⁴</p>
<p>Low birthweight</p>	<p>The World Health Organization defines low birthweight as weight at birth less than 2,500 grams (5.5 pounds).</p> <p>Low birthweight can be caused by intrauterine growth restriction as well as prematurity. It is associated with fetal and neonatal morbidity and mortality as well as impaired growth and cognitive development in childhood.⁸⁵ It is also associated with chronic disease in adults, such cardiovascular disease, type 2 diabetes and metabolic syndrome.⁸⁶</p> <p>Many environmental hazards are associated with low birthweight, including air pollution, tobacco smoke and pesticides.⁸⁷</p>
<p>Birth defects</p>	<p>Air pollution, consumption of foods contaminated with fumonisins, extreme heat, hazardous waste and solvents are just some of the exposures that have been shown to be linked to various birth defects.</p> <p>These defects include malformation of the central nervous system, heart, abdominal organs, limbs and other structures.⁸⁸</p>
<p>Numerous chronic conditions in childhood and adulthood</p>	<p>Beyond birth, diseases related to antenatal exposures can develop in childhood, adolescence and adulthood.</p> <p>Antenatal exposure to air pollution is associated with chronic childhood conditions, such as asthma, allergic disease, neurodevelopmental disorders, obesity and high blood pressure. It is also linked to adulthood diseases such as cardiovascular disease and chronic obstructive pulmonary disease, the third-leading cause of death worldwide.⁸⁹</p> <p>Antenatal pesticide exposure is associated with neurodevelopmental disorders and childhood cancers including neuroblastoma and leukaemia, the most common childhood cancer.⁹⁰</p> <p>Exposure to mercury during pregnancy can lead to devastating neurological outcomes in a child.⁹¹</p>

References

- 1 Etzel, Ruth A., and Philip J. Landrigan, 'Children's Exquisite Vulnerability to Environmental Exposures', ch. 2 in *Textbook of Children's Environmental Health*, edited by Ruth A. Etzel and Philip J. Landrigan, Oxford University Press, New York, 2024, pp. 18–27.
- 2 Yao, Xiaoxi, et al., 'Fertility Loss: Negative effects of environmental toxicants on oogenesis', *Frontiers in Physiology*, vol. 14, 4 August 2023, e.1219045; Greeson, Katherine W., et al., 'Inheritance of Paternal Lifestyles and Exposures Through Sperm DNA Methylation', *Nature Reviews Urology*, vol. 20, no. 6, 18 January 2023, pp. 356–370.
- 3 Stephenson, Judith, et al., 'Before the Beginning: Nutrition and lifestyle in the preconception period and its importance for future health', *Lancet*, vol. 391, no. 10132, 5 May 2018, pp. 1830–1841.
- 4 Perera, Frederica, and Julie Herbstman, 'Prenatal Environmental Exposures, Epigenetics, and Disease', *Reproductive Toxicology*, vol. 31, no. 3, 12 September 2011, pp. 363–373.
- 5 Ibid.; Li, Yu-Fen, et al., 'Maternal and Grandmaternal Smoking Patterns are Associated with Early Childhood Asthma', *Chest*, vol. 127, no. 4, April 2005, pp. 1232–1241.
- 6 Agency for Toxic Substances and Disease Registry, 'How Can Parents' Preconception Exposures and In Utero Exposures Affect a Developing Child?' <https://archive.cdc.gov/www_atsdr_cdc_gov/csem/pediatric-environmental-health/preconception.html>, accessed 5 December 2024.
- 7 Yao et al., 'Fertility Loss'; Greeson et al., 'Inheritance of Paternal Lifestyles and Exposures'; Moore, Keith L., T. Vidhya Persaud and Mark G. Torchia, *The Developing Human: Clinically oriented embryology*, 11th ed., Elsevier, London, 2020.
- 8 Nilsson, Eric E., Ingrid Sadler-Riggelman and Michael K. Skinner, 'Environmentally Induced Epigenetic Transgenerational Inheritance of Disease', *Environmental Epigenetics*, vol. 4, no. 2., April 2018.
- 9 Skinner, Michael K., et al., 'Ancestral Dichlorodiphenyltrichloroethane (DDT) Exposure Promotes Epigenetic Transgenerational Inheritance of Obesity', *BMC Medicine*, vol. 11, no. 228, October 2013.
- 10 United States Centers for Disease Control and Prevention, 'Epigenetics, Health and Disease', 13 November 2024, <www.cdc.gov/genomics-and-health/epigenetics/index.html>, accessed 5 December 2024.
- 11 Perera and Herbstman, 'Prenatal Environmental Exposures'.
- 12 World Health Organization, 'Arsenic', 7 December 2022, <www.who.int/news-room/fact-sheets/detail/arsenic>, accessed 11 September 2024.
- 13 Ortiz-Garcia, Nancy Y., et al., 'Maternal Exposure to Arsenic and its Impact on Maternal and Fetal Health: A review', *The Cureus Journal of Medical Science*, vol. 15, no. 11, 21 November 2023, e49177.
- 14 Tolins, Molly, Mathuros Ruchirawat and Philip Landrigan, 'The Developmental Neurotoxicity of Arsenic: Cognitive and behavioral consequences of early life exposure', *Annals of Global Health*, vol. 80, no. 4, July 2014, pp. 303–314.
- 15 Song, Yongye, et al., 'A Study of the Fluid Intake, Hydration Status, and Health Effects among Pregnant Women in Their Second Trimester in China: A cross-sectional study', *Nutrients*, vol. 15, no. 7, 2 April 2023, e1739.
- 16 Tran, Nguyen-Toan, et al., 'Cholera in Pregnancy: A systematic review and meta-analysis of fetal, neonatal, and maternal mortality', *PLOS One*, vol. 10, no. 7, 15 July 2015, e0132920; El Hayek, Pamala, et al., 'Cholera Infection Risks and Cholera Vaccine Safety in Pregnancy', *Infectious Diseases in Obstetrics and Gynecology*, 22 May 2023, e4563797.
- 17 Bonell, Ana, et al., 'Effect of Heat Stress in the First 1000 Days of Life on Fetal and Infant Growth: A secondary analysis of the ENID randomised controlled trial', *Lancet Planetary Health*, vol. 8, no. 10, October 2024, pp. e734–e743.
- 18 Augusto, Ana Lucia Pires, et al., 'Household Food Insecurity Associated with Gestational and Neonatal Outcomes: A systematic review', *BMC Pregnancy and Childbirth*, vol. 20, no. 229, 17 April 2020; Carmichael, Suzan L., et al., 'Maternal Food Insecurity is Associated with Increased Risk of Certain Birth Defects', *Journal of Nutrition*, vol. 137, no. 9, September 2007, pp. 2087–2092; Richterman, Aaron M., et al., 'Food Insecurity as a Risk Factor for Preterm Birth: A prospective facility-based cohort study in rural Haiti', *BMJ Global Health*, vol. 5, no. 7, July 2020, e002341.
- 19 Augusto et al., 'Household Food Insecurity'.
- 20 Du, Rebecca Y., et al., 'Systematic Review of Clinician Awareness of Mycotoxin Impact in Neural Tube Defects and Best Practices for Pediatric Neurosurgeons: Implications for public health and policy', *Child's Nervous System*, vol. 35, no. 4, April 2019, pp. 637–644; American Academy of Pediatrics Council on Environmental Health, 'Birth Defects and Other Adverse Developmental Outcomes', in *Pediatric Environmental Health*, 4th ed., edited by Ruth A. Etzel and Sophie J. Balk, American Academy of Pediatrics, Itasca, Ill., 2019, p. 862.
- 21 LoMauro, Antonella, and Andrea Aliverti, 'Respiratory Physiology of Pregnancy: Physiology masterclass', *Breathe*, vol. 11, no. 4, December 2015, pp. 297–301.
- 22 Saenen, Nelly D., et al., 'Air Pollution-Induced Placental Alterations: An interplay of oxidative stress, epigenetics, and the aging phenotype?', *Clinical Epigenetics*, vol. 11, no. 124, 17 September 2019; Johnson, Natalie M., et al., 'Air Pollution and Children's Health: A review of adverse effects associated with prenatal exposure from fine to ultrafine particulate matter', *Environmental Health and Preventative Medicine*, vol. 26, no. 72, 12 July 2021.
- 23 Nyadanu, Sylvester D., et al., 'Prenatal Exposure to Ambient Air Pollution and Adverse Birth Outcomes: An umbrella review of 36 systematic reviews and meta-analyses', *Environmental Pollution*, vol. 306, 1 August 2022, e119465.
- 24 Ghosh, Rakesh, et al., 'Ambient and Household PM2.5 Pollution and Adverse Perinatal Outcomes: A meta-regression and analysis of attributable global burden for 204 countries and territories', *PLOS Medicine*, vol. 18, no. 9, 28 September 2021, e1003718.
- 25 Nyadanu et al., 'Prenatal Exposure to Ambient Air Pollution'.
- 26 Zhang, Huanhuan, et al., 'Ambient Air Pollution and Stillbirth: An updated systematic review and meta-analysis of epidemiological studies', *Environmental Pollution*, vol. 278, 1 June 2021, e116752.
- 27 Lin, Li-Zi, et al., 'The Epidemiological Evidence Linking Exposure to Ambient Particulate Matter with Neurodevelopmental Disorders: A systematic review and meta-analysis', *Environmental Research*, vol. 209, June 2022, e112876.
- 28 Hazlehurst, Marnie F., et al., 'Maternal Exposure to PM(2.5) During Pregnancy and Asthma Risk in Early Childhood: Consideration of phases of fetal lung development', *Environmental Epidemiology*, vol. 5, no. 2, 2 March 2021, e130.
- 29 Perera, Frederica, and Kari Nadeau, 'Climate Change, Fossil-Fuel Pollution, and Children's Health', *New England Journal of Medicine*, vol. 386, no. 24, 15 June 2022, pp. 2303–2314.
- 30 United States Centers for Disease Control and Prevention, 'About Secondhand Smoke', 15 May 2024, <www.cdc.gov/tobacco/second-hand-smoke/index.html>, accessed 28 November 2024.
- 31 Zhou, Sherry, et al., 'Physical, Behavioral, and Cognitive Effects of Prenatal Tobacco and Postnatal Secondhand Smoke Exposure', *Current Problems in Pediatric and Adolescent Health Care*, vol. 44, no. 8, September 2014, pp. 219–241.

- 32 World Health Organization, 'Tobacco Control to Improve Child Health and Development: Thematic brief', WHO, Geneva, 2021.
- 33 Maciag, Michelle C., Aroub Yousuf and Marissa Hauptman, 'Impact of Prenatal Exposure to Smoking on Child Health', *Clinical Obstetrics and Gynecology*, vol. 65, no. 2, June 2022, pp. 388–396.
- 34 Hussain, Tarique, et al., 'The Role of Oxidative Stress and Antioxidant Balance in Pregnancy', *Mediators of Inflammation*, 27 September 2021, e9962860.
- 35 Gitto, Eloisa, et al., 'Oxidative Stress of the Newborn in the Pre- and Postnatal Period and the Clinical Utility of Melatonin', *Journal of Pineal Research*, vol. 46, no. 2, March 2009, pp. 128–139.
- 36 Mathiesen, Line, et al., 'Fetal Exposure to Environmental Chemicals; Insights from placental perfusion studies', *Placenta*, vol. 106, March 2021, pp. 58–66; Dai, Yifeng, et al., 'Early-Life Exposure to Widespread Environmental Toxicants and Maternal-Fetal Health Risk: A focus on metabolomic biomarkers', *Science of the Total Environment*, vol. 739, 15 October 2020, e139626.
- 37 Gitto et al., 'Oxidative Stress of the Newborn'.
- 38 Trivett, Cara, Zoe J. Lees and Dilys J. Freeman, 'Adipose Tissue Function in Healthy Pregnancy, Gestational Diabetes Mellitus and Pre-eclampsia', *European Journal of Clinical Nutrition*, vol. 75, no. 12, 15 June 2021, pp. 1745–1756.
- 39 Kapraun, Dustin F., et al., 'A Generic Pharmacokinetic Model for Quantifying Mother-to-Offspring Transfer of Lipophilic Persistent Environmental Chemicals', *Toxicological Sciences*, vol. 189, no. 2, 24 September 2022, pp. 155–174.
- 40 Mustieles, Vicente, and Juan P. Arrebola, 'How Polluted is Your Fat? What the study of adipose tissue can contribute to environmental epidemiology', *Journal of Epidemiology and Community Health*, vol. 74, no. 5, May 2020, pp. 401–407.
- 41 Merrill, Alyssa K., Marisia Sobolewski and Marsha Susiarjo, 'Exposure to Endocrine Disrupting Chemicals Impacts Immunological and Metabolic Status of Women During Pregnancy', *Molecular and Cellular Endocrinology*, vol. 577, 1 November 2023, e.112031.
- 42 Sanghavi, Monika, and John D. Rutherford, 'Cardiovascular Physiology of Pregnancy', *Circulation*, vol. 130, no. 12, 16 September 2014, pp. 1003–1008.
- 43 Aguree, Sixtus, and Alison D. Gernand, 'Plasma Volume Expansion Across Healthy Pregnancy: A systematic review and meta-analysis of longitudinal studies', *BMC Pregnancy and Childbirth*, vol. 19, no. 508, 2019.
- 44 Camaschella, Clara, 'Iron Deficiency', *Blood*, vol. 133, no. 1, 3 January 2019, pp. 30–39.
- 45 Wyrwoll, Catilin S., 'Rising Stars: The heat is on – How does heat exposure cause pregnancy complications?', *Journal of Endocrinology*, vol. 259, no. 1, 25 August 2023, e230030.
- 46 Ibid.
- 47 Lakhoo, Darshnika P., et al., 'A Systematic Review and Meta-analysis of Heat Exposure Impacts on Maternal, Fetal and Neonatal Health', *Nature Medicine*, 5 November 2024, print publication forthcoming.
- 48 Chan, Monica Y., and Mary Alice Smith, 'Infections in Pregnancy', vol. 5 ch. 16 in *Comprehensive Toxicology*, 3rd ed., edited by Charlene A. McQueen, Elsevier, Amsterdam, 2018, pp. 232–249.
- 49 Ibid.
- 50 Ibid.
- 51 Ibid.
- 52 Rogerson, Steven J., and Holger W. Unger, 'Pregnancy and Malaria: The perfect storm', *Current Opinion in Infectious Diseases*, vol. 35, no. 5, October 2022, pp. 410–416.
- 53 Zorrilla, Carmen D., et al., 'Zika Virus Infection in Pregnancy: Maternal, fetal, and neonatal considerations', *Journal of Infectious Diseases*, vol. 216, suppl. 10, 15 December 2017, pp. S891–S896; World Health Organization, 'Zika Virus', 8 December 2022, <www.who.int/news-room/fact-sheets/detail/zika-virus>, accessed 5 December 2024.
- 54 Dahiya, Nisha, et al., 'ZIKV: Epidemiology, infection mechanism and current therapeutics', *Frontiers in Tropical Diseases*, vol. 3, 23 January 2023, 1059283.
- 55 Krauer, Fabienne, et al., 'Zika Virus Infection as a Cause of Congenital Brain Abnormalities and Guillain-Barré Syndrome: Systematic review', *PLOS Medicine*, vol. 14, no. 1, 3 January 2017, e1002203; Musso, Didier, Albert I. Ko and David Baud, 'Zika Virus Infection: After the pandemic', *New England Journal of Medicine*, vol. 381, no. 15, 4 October 2019, pp. 1444–1457.
- 56 Marbán-Castro, Elana, et al., 'Zika Virus Infection in Pregnant Women and Their Children: A review', *European Journal of Obstetrics & Gynecology and Reproductive Biology*, vol. 265, October 2021, pp. 162–168.
- 57 Burton, Graham J., and Abigail L. Fowden, 'The Placenta: A multifaceted, transient organ', *Philosophical Transactions of the Royal Society of London*, vol. 370, no. 1663, 5 March 2015, e20140066.
- 58 Khorami-Sarvestani, Sara, et al., 'Placenta: An old organ with new functions', *Frontiers in Immunology*, vol. 15, 19 April 2024, e1385762.
- 59 Mathiesen et al., 'Fetal Exposure to Environmental Chemicals': Dai et al., 'Early-Life Exposure to Widespread Environmental Toxicants'.
- 60 Ortiz-Garcia et al., 'Maternal Exposure to Arsenic'.
- 61 Megli, Christina J., and Carolyn B. Coyne, 'Infections at the Maternal–Fetal Interface: An overview of pathogenesis and defence', *Nature Reviews Microbiology*, vol. 20, no. 2, February 2022, pp. 67–82.
- 62 Gingrich, Jeremy, Elvis Ticiani and Alumdena Veiga-Lopez, 'Placenta Disrupted: Endocrine disrupting chemicals and pregnancy', *Trends in Endocrinology & Metabolism*, vol. 31, no. 7, July 2020, pp. 508–524.
- 63 Bloom, Michael, Meghana Varde and Roger Newman, 'Environmental Toxicants and Placental Function', *Best Practice and Research in Clinical Obstetrics and Gynaecology*, vol. 85, part B, December 2022, pp. 105–120.
- 64 Dimitriadis, Evdokia, et al., 'Pre-eclampsia', *Nature Reviews Disease Primers*, vol. 9, art. 8, 16 February 2023.
- 65 Ibid.
- 66 Mazumder, Hoimonty, et al., 'Maternal Health Outcomes Associated with Ambient Air Pollution: An umbrella review of systematic reviews and meta-analyses', *Science of the Total Environment*, vol. 914, 1 March 2024, 169792.
- 67 Moore, Persaud and Torchia, *The Developing Human*.
- 68 Breton-Larrivée, Mélanie, Elizabeth Elder and Serge McGraw, 'DNA Methylation, Environmental Exposures and Early Embryo Development', *Animal Reproduction*, vol. 16, no. 3, 23 October 2019, pp. 465–474.
- 69 Isaevska, Elena, et al., 'Exposure to Ambient Air Pollution in the First 1000 Days of Life and Alterations in the DNA Methylome and Telomere Length in Children: A systematic review', *Environmental Research*, vol. 193, February 2021, e110504.
- 70 Moore, Persaud and Torchia, *The Developing Human*.
- 71 Decrue, Fabienne, et al., 'Increased Impact of Air Pollution on Lung Function in Preterm versus Term Infants: The BILD study', *American Journal of Respiratory and Critical Care Medicine*, vol. 205, no. 1, 1 January 2022, pp. 99–107.
- 72 American College of Obstetricians and Gynecologists, 'How Your Fetus Grows During Pregnancy', January 2024, <www.acog.org/womens-health/faqs/how-your-fetus-grows-during-pregnancy>, accessed 23 August 2024.

- 73 Moore, Persaud and Torchia, *The Developing Human*.
- 74 Di Renzo, Gian C., et al., 'International Federation of Gynecology and Obstetrics Opinion on Reproductive Health Impacts of Exposure to Toxic Environmental Chemicals', *International Journal of Gynecology & Obstetrics*, vol. 131, no. 3, December 2015, pp. 219–225.
- 75 World Health Organization, 'Household Air Pollution', 16 October 2024, <www.who.int/news-room/fact-sheets/detail/household-air-pollution-and-health>, accessed 5 December 2024.
- 76 World Health Organization, *Children and Digital Dumpsites: E-waste exposure and child health*, WHO, Geneva, 2021; Perks, Rachel, 'Let's Support Women in Artisanal and Small-Scale Mining', World Bank Blogs, Washington, D.C., 8 March 2024, <<https://blogs.worldbank.org/en/energy/lets-support-women-artisanal-and-small-scale-mining>>, accessed 12 December 2024; Food and Agriculture Organization of the United Nations and Rotterdam Convention, *Addressing Gender Issues in Pesticide Management*, FAO, Rome, 2022.
- 77 Gender and Hazardous Substances: Report of the Special Rapporteur on the implications for human rights of the environmentally sound management and disposal of hazardous substances and wastes, Marcos Orellana, A/79/163, United Nations, New York, 16 July 2024; Strategic Approach to International Chemicals Management and the International Pollutants Elimination Network, *Women, Chemicals and the SDGs*, IPEN, Geneva, 2022.
- 78 McNab, Shanon, and Lynn Freedman, *Maternal Newborn Health and the Urban Poor: A global scoping*, Columbia University Mailman School of Public Health, New York, 2016.
- 79 Yao, Xiaodie, et al., 'Environmental Pollutants Exposure and Gestational Diabetes Mellitus: Evidence from epidemiological and experimental studies', *Chemosphere*, vol. 332, August 2023, e138866.
- 80 Mazumder et al., 'Maternal Health Outcomes'.
- 81 Nyadanu, et al., 'Prenatal Exposure to Ambient Air Pollution'; Zhang et al., 'Ambient Air Pollution and Stillbirth'; Anderson, Mark E., Daniel C. Johnson and Holly A. Batal, 'Sudden Infant Death Syndrome and Prenatal Maternal Smoking: Rising attributed risk in the Back to Sleep era', *BMC Medicine*, vol. 3, no. 1, art. 4, 11 January 2005.
- 82 Etzel, Ruth A., 'Is the Environment Associated With Preterm Birth?', *JAMA Network Open*, vol. 3, no. 4, 2020, e202239.
- 83 Ibid.
- 84 World Health Organization, 'Preterm Birth', 10 May 2023, <www.who.int/news-room/fact-sheets/detail/preterm-birth>, accessed 26 May 2023.
- 85 World Health Organization, 'Low Birth Weight', <<https://www.who.int/data/nutrition/nlis/info/low-birth-weight>>, accessed 5 December 2024.
- 86 Smith, C. J., et al., 'The Impact of Birth Weight on Cardiovascular Disease Risk in the Women's Health Initiative', *Nutrition, Metabolism & Cardiovascular Diseases*, vol. 26, no. 3, March 2016, pp. 239–245.
- 87 Triche, Elizabeth W., and Nazli Hossain, 'Environmental Factors Implicated in the Causation of Adverse Pregnancy Outcome', *Seminars in Perinatology*, vol. 31, no. 4, August 2007, pp. 240–242.
- 88 'Birth Defects and Other Adverse Developmental Outcomes', p. 865.
- 89 Núñez-Sánchez, María Á., et al., 'Inherited Epigenetic Hallmarks of Childhood Obesity Derived from Prenatal Exposure to Obesogens', *International Journal of Environmental Research and Public Health*, vol. 20, no. 6, 7 March 2023, e4711; Hazlehurst et al., 'Maternal Exposure to PM(2.5) During Pregnancy'; Brumberg, Heather L., et al., 'Ambient Air Pollution: Health hazards to children', *Pediatrics*, vol. 147, no. 6, June 2021, e2021051484; Tingskov Pedersen, Casper E., et al., 'Prenatal Exposure to Ambient Air Pollution is Associated with Early Life Immune Perturbations', *Journal of Allergy and Clinical Immunology*, vol. 151, no. 1, January 2023, pp. 212–221; Kim, Juyong B., et al., 'Cumulative Lifetime Burden of Cardiovascular Disease From Early Exposure to Air Pollution', *Journal of the American Heart Association*, vol. 9, no. 6, 17 March 2020, e014944; Deolmi, Michela, et al., 'Early Origins of Chronic Obstructive Pulmonary Disease: Prenatal and early life risk factors', *International Journal of Environmental Research and Public Health*, vol. 20, no. 3, February 2023, e2294.
- 90 Hertz-Picciotto, Irva, et al., 'Organophosphate Exposures During Pregnancy and Child Neurodevelopment: Recommendations for essential policy reforms', *PLOS Medicine*, vol. 15, no. 10, October 2018, e1002671; Van Maele-Fabry, Geneviève, Laurence Gamet-Payraastre and Dominique Lison, 'Household Exposure to Pesticides and Risk of Leukemia in Children and Adolescents: Updated systematic review and meta-analysis', *International Journal of Hygiene and Environmental Health*, vol. 222, no. 1, January 2019, pp. 49–67; Khan, Aiza, Joseph Feulefack and Consolato M. Sergi, 'Pre-conceptional and Prenatal Exposure to Pesticides and Pediatric Neuroblastoma: A meta-analysis of nine studies', *Environmental Toxicology and Pharmacology*, vol. 90, February 2022, 103790.
- 91 World Health Organization, 'Preventing Disease Through Health Environments: Exposure to mercury – A major public health concern', 2nd ed., WHO, Geneva, 8 April 2021.



© UNICEF/UN0731260/zquierdo

Acknowledgements

Author: Maria Brown, with guidance from Abheet Solomon

UNICEF reviewers: Maaïke Arts, Alexandre Boon, Ulrike Gilbert, Katelyn Greer, Tomomi Kitamura, Desiree Narvaez, Chemba Raghavan, Eduardo Garcia Rolland, Kam Sripada

External reviewers: Perna Banati, Ruth Etzel, Julia Gorman, Phil Landrigan, Frederica Perera, Skye Wheeler

The statements in this document should not be taken as representing the official position and policies of UNICEF.

The document will be revised as new information and contributions becomes available. Suggestions and comments are welcome and may be sent to ceh@unicef.org.

United Nations Children's Fund

3 United Nations Plaza
New York, NY 10017, USA
www.unicef.org

© United Nations Children's Fund (UNICEF), December 2024

unicef 
for every child